




Biology Lab Activities: Monocot and Dicot Plant Structure

Adrienne Devlin, Executive Editor at Visible Body

How to use this manual

This lab manual is intended for use with the <u>Visible Biology</u> product.

Where to find 3D models



How to save answers

1. Have Adobe Reader installed on your computer.


Windows: https://get.adobe.com/reader/

Mac: https://helpx.adobe.com/acrobat/kb/install-reader-dc-mac-os.html

2. Download each lab file to your computer.



- 3. Open the downloaded file in Adobe Reader.
 Right-click on the file. In the menu that appears, go to "Open with..." and select Adobe Reader from the submenu.
- 4. Type your answers into the boxes to complete the lab and select the "Save" icon to save the lab.



5. Submit your saved version of the lab to your instructor via email, dropbox, Google Drive, or however your instructor has requested.

Name:	Date:

Biology Lab Activities: Monocot and Dicot Plant Structure

Background Questions

Read through the Monocot and Dicot Overview article on the Visible Body Biology Learn Site (https://www.visiblebodv.com/learn/biologv/monocot-dicot/overview).

Based on what you've learned in class, in your textbook, from the Biology Learn Site article, and from using Visible Biology, answer the following questions about the structures that make up flowering plants.

- 1. What are the three main parts of plants?
- 2. There are three types of tissue found in each part of the plant: dermal, vascular, and ground tissue. Answer the following questions about the structure and functions of these tissues.
 - a. Dermal tissue is composed of a single layer of ______, and it forms the outer tissue layer of the plant. b. What are the two main functions of dermal tissue?
 - c. Vascular tissue is primarily composed of _____ and ____, which form vascular bundles throughout the plant's interior.
 - d. What is the main function of vascular tissue?
 - e. Ground tissue can be composed of three types of cells: _____, and _____. It makes up the majority of the plant's interior.
 - f. What are the two main functions of ground tissue?
 - i.

i. ii.

ii.


3.	There	are two main types of flowering plan	its, monocots and	dicots (also known as
	eudico	ots). Complete the following sentenc	es to compare the	se two plant types.
	a.	Monocots usually have	ereas dicots usually have	
		cotyledon(s). This structure	pecause it supplies	
		to the	e plant embryo.	
	b.	The leaves of monocots are usuall	у	, with
		veins. On the otl	es of dicots are more varied	
		in shape and size, usually with		veins.
	C.	In monocots, the vascular bundles	in the stem are	, whereas in
		dicots, they are arranged in a		
	d.	Monocots have flower parts in grou	ups of	, whereas dicots have
		flower parts in groups of		
	e.	Monocots are monosulcate, meani	ng their	
		Dicots are trisulcate, meaning their	,	

Lab 1: Monocot and Dicot Plant Roots

Activity 1: Label a monocot plant root

- 1. Launch the view
 - Launch Visible Biology.
 - Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 1. Monocot Root.
- 2. Label the image below
 - Explore the 3D model of the monocot root to find the structures you need to label.
 - Fill in the blanks to label the structures from the list below.

Word List:			
Cortex	 Pericycle	 Root hair	
Endodermis	 Phloem	 Stele	
Epidermis	 Pith	 Xylem	




Lab 1: Monocot and Dicot Plant Roots

Activity 2: Label a dicot plant root

- 1. Launch the view
 - Launch Visible Biology.
 - Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 2. Dicot Root.
- 2. Label the image below
 - Explore the 3D model of the dicot root to find the structures you need to label.
 - o Fill in the blanks to label the structures from the list below.

Word List:

Cambium	 Epidermis	 Stele
Connective tissue (parenchyma)	 Pericycle	 Xylem
Cortex	 Phloem	
Endodermis	 Root hair	



Name:	Date:

Lab 1: Monocot and Dicot Plant Roots

Activity 3: Compare monocot and dicot plant roots

As you've learned from labeling monocot and dicot plant roots in Activities 1 and 2, they share most of the same structures, but those structures are arranged differently within each type of root. Refer to your labeled images and the content in Visible Biology to answer the following questions, comparing monocot and dicot plant roots.

	tefer to your labeled images and the content in Visible Biology to answer the following ons, comparing monocot and dicot plant roots.
1.	Based on your labeled monocot and dicot roots, match the following root structures with their descriptions. a. Root hairs b. Endodermis c. Pericycle d. Epidermis e. Stele
	A single layer of parenchyma cells that forms the boundary between the cortex and the stele The central part of the root where the xylem and phloem develop The protective outer layer that prevents damage to the root Structures that enhance the root's total surface area to maximize water and mineral absorption The outermost layer of the stele, composed of parenchyma cells that can divide and give rise to lateral roots
2.	How would you describe the overall shape and structure of monocot vs. dicot roots, in 2–3 sentences?
3.	The root's ground tissue can be found in a region called the In this region, monocot and dicot roots both have, but monocot roots can also have
4.	Monocot and dicot roots both have vascular cylinders. Complete the following sentences about how the vascular structures differ in monocot and dicot roots.


a.	roots have a larger number of vascular structures than				
	roots.				
b.	In monocot roots, the vascular structures are arranged in a				
	between the outer cortex and the central				
C.	In dicot roots, the vascular structures are located in the of the				
	root. Dicot steles contain an additional component called, which	ch			
	is located between the xylem and phloem. The meristem cells within this				
	structure facilitate secondary growth, dividing to create new xylem and phloem				
	cells, and thus widening the root's girth.				

Lab 2: Monocot and Dicot Plant Stems

Activity 1: Label a monocot plant stem

- 1. Launch the view
 - Launch Visible Biology.
 - o Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 3. Monocot Stem.
- 2. Label the image below
 - o Explore the 3D model of the monocot stem to find the structures you need to label.
 - Fill in the blanks to label the structures from the list below.

<u>vvora List:</u>			
Epidermis	 Phloem	 Xylem	
Ground tissue	 Sclerenchyma		
Hypodermis	Vascular bundle		




Lab 2: Monocot and Dicot Plant Stems

Activity 2: Label a dicot plant stem

- 1. Launch the view
 - Launch Visible Biology.
 - o Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 4. Dicot Stem.
- 2. Label the image below
 - o Explore the 3D model of the dicot stem to find the structures you need to label.
 - o Fill in the blanks to label the structures from the list below.

Word List:			
Cambium	 Pith	 Vascular bundle	
Cortex	 Phloem	 Xylem	
Epidermis	Sclerenchyma		



Lab 2: Monocot and Dicot Plant Stems

Activity 3: Compare monocot and dicot plant stems

As you've learned from labeling monocot and dicot plant stems in Activities 1 and 2, they share several of the same structures, but those structures are arranged differently within each type of stem. Refer to your labeled images and the content in Visible Biology to answer the following questions, comparing monocot and dicot plant stems.

	,	
1.	sclerer arrang	ajority of the stem is composed of ground tissue, which can include parenchyma, nchyma, and collenchyma cells. In most ground tissue, the cells are loosely ed and there is space between them. Briefly explain why this space is important to action of the plant stem.
2.	vascul stem's	can see in your labeled monocot and dicot stems, they both contain several ar bundles. However, these vascular structures are arranged differently within the ground tissue. Complete the following sentences about the structure and location se vascular bundles.
	a.	The interior of the vascular bundles is composed of and the exterior is composed of Dicot stems have an additional
		component called, which is located between the xylem and phloem. The meristem cells within this structure facilitate secondary growth, dividing to create new xylem and phloem cells, and thus widening the stem's girth.
	b.	Monocot stems have vascular bundles that are
	C.	Dicot stems have vascular bundles that form a, separating the

stem's ground tissue into an outer _____ and a central _____.


Lab 3: Monocot and Dicot Plant Leaves

Activity 1: Label a monocot plant leaf

- 1. Launch the view
 - Launch Visible Biology.
 - o Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 5. Monocot Leaf.
- 2. Label the image below
 - o Explore the 3D model of the monocot leaf to find the structures you need to label.
 - o Fill in the blanks to label the structures from the list below.

Word List:

Bulliform cell	 Lower epidermis	 Sclerenchyma	
Bundle sheath	Mesophyll	Stoma	
Cuticle	Parenchyma	Upper epidermis	
Guard cell	Phloem	Xylem	




Lab 3: Monocot and Dicot Plant Leaves

Activity 2: Label a dicot plant leaf

- 1. Launch the view
 - Launch Visible Biology.
 - o Navigate to Study/Lab Activities, and find the Monocot and Dicot Plant Structure Lab section.
 - Select view 6. Dicot Leaf.
- 2. Label the image below
 - Explore the 3D model of the dicot leaf to find the structures you need to label.
 - o Fill in the blanks to label the structures from the list below.

Word List:

Bundle sheath Collenchyma Cuticle Guard cell	 Palisade mesophyll Phloem Spongy mesophyll Stoma	_ _ _	Upper epidermis Vascular bundle Xylem	
Lower epidermis	 Trichome			



Lab 3: Monocot and Dicot Plant Leaves

Activity 3: Compare monocot and dicot plant leaves

As you've learned from labeling monocot and dicot plant leaves in Activities 1 and 2, they share many of the same structures, but those structures are arranged differently within each type of leaf. Refer to your labeled images and the content in Visible Biology to answer the following questions, comparing monocot and dicot plant leaves.

1.	a. b.	The leaf's dermal cells secrete a waxy substance that forms the This structure forms a protective covering and helps the leaf The leaf's epidermis has several small pores, called, which facilitate and allow water vapor to exit the leaf. Each pore is surrounded by two, which open and close the pore. In, stomata are found on the upper and lower surfaces of the leaves. In, most stomata are on the bottom surface. This allows dicot leaves to while having most stomata open.
2.	a.	lete the following sentences on the ground tissue of leaves. The leaf's ground tissue, called, separates the upper and lower The ground tissue is mostly composed of parenchyma cells that carry out
	D.	within the leaf.
	C.	Unlike most monocot leaves, which have only one type of mesophyll, the ground tissue of dicot leaves is often arranged into two distinct types, the and the
3.	As you've probably noticed in your labeled monocot and dicot leaves, the vascular structures form parallel veins in monocot leaves and net-like veins in dicot leaves. The xylem and phloem in these veins are connected to the stem's vascular structures. In 2–3 sentences, explain why this structure is important to the functions of the leaf and the res of the plant.	

Name:	Date:
Name:	D

Lab 4: Dermal, Vascular, and Ground Tissue in Plants

Activity 1: Dermal Tissue

As you've learned from labeling monocot and dicot plant roots, stems, and leaves in Labs 1–3, all these plant parts are composed of dermal, vascular, and ground tissues. Refer to your labeled images, as well as the 3D models in Visible Biology, to answer the following questions on

the	dermal tissue of plants.
1.	What dermal tissue structure is present in roots, stems, and leaves?
2.	How would you describe the basic structure of this dermal tissue?
3.	Although this dermal structure is shared by all three plant parts, what is unique about its structure in leaves?
4.	Match each of the following dermal tissue structures with its function. a. Cuticle b. Root hair c. Stoma d. Guard cell
	Allows gases and water vapor to move between the interior plant structures and the surrounding air Helps the plant retain water and prevents pathogens from entering the plant Opens and closes a pore Absorbs water and minerals from the ground

Lab 4: Dermal, Vascular, and Ground Tissue in Plants

Activity 2: Vascular Tissue

As you've learned from labeling monocot and dicot plant roots, stems, and leaves in Labs 1–3, all these plant parts are composed of dermal, vascular, and ground tissues. Refer to your labeled images, as well as the 3D models in Visible Biology, to answer the following questions on the vascular tissue of plants.

vaooaic	ai lissue di piants.		
	• •	ascular tissues that are present	in roots, stems, and
•	•		
a.	The carrie the other plant parts.	s water and dissolved minerals	from the to
b.	c compounds from the		
C.		•	at maturity.
senter	nces to compare the arrang	•	•
a.			
b.		<u> </u>	
		In monocot stems, they are	;
		In dicot stems, they	form a
		between the outer	and the inner
C.	•		
		veins. In monocot leaves, the	e veiris are
	What seleaves Comp a. b. c. Vascusenter stems a. b.	What are the two main types of valeaves? Complete the following sentences a. The carrie the other plant parts. b. The carrie to the other of the other carrie to the other	What are the two main types of vascular tissues that are present leaves? Complete the following sentences about the main functions of the a. The carries water and dissolved minerals the other plant parts. b. The carries dissolved sugars and organized to the other plant parts. c. Xylem cells have walls, and they are Phloem cells have walls, and they are plant. Vascular tissue is arranged differently in each part of the plant. Consentences to compare the arrangement and structure of the vascular tissue is arranged differently in each part of the plant.

4. How would you explain the cambium in 2–3 sentences?

Name:	Date:
-------	-------

Lab 4: Dermal, Vascular, and Ground Tissue in Plants

Activity 3: Ground Tissue

As you've learned from labeling monocot and dicot plant roots, stems, and leaves in Labs 1–3, all these plant parts are composed of dermal, vascular, and ground tissues. Refer to your labe on

	•	es, as well as the 3D mo d tissue of plants.	dels in Visible Biology, to answer the following questions		
1.	What	are the three main types	of ground tissue cells?		
1.	Comp cell.	Complete the following sentences about the main functions of each type of ground tissue cell.			
	a.	The most abundant typ	e, cells, perform vital plant functions		
			nesis, respiration, gas exchange, and water and starch		
	b.		cells provide elastic support that facilitates stem and leaf		
		growth.			
	C.		cells provide inelastic support that maintains the rigid		
		structure of plant parts	that are done growing.		
2	Which	narts of roots, stems, a	nd leaves are composed of ground tissue?		